

Effects of loud noise exposure on DNA integrity in rat adrenal gland

Giada Frenzilli, Paola Lenzi, Vittoria Scarcelli, Francesco Fornai, Antonio Pellegrini, Paola Soldani, Antonio Paparelli, and Marco Nigro

doi:10.1289/ehp.7249 (available at http://dx.doi.org/)
Online 22 September 2004



Effects of loud noise exposure on DNA integrity in rat adrenal gland.

Giada Frenzilli, Paola Lenzi, Vittoria Scarcelli, Francesco Fornai, Antonio Pellegrini, Paola Soldani, Antonio Paparelli, Marco Nigro.

Dipartimento di Morfologia Umana e Biologia Applicata, University of Pisa, Via Roma, 55 56126 Pisa, Italy.

Address correspondence to: Giada Frenzilli, Dipartimento di Morfologia Umana e Biologia Applicata, Sez. Biologia e Genetica University of Pisa, Via Volta, 4 - 56126 Pisa. Telephone +39 050 2219111. Fax +39 050 2219101- E-Mail: giada@biomed.unipi.it.

Running title: Noise-induced DNA damage in the rat adrenal gland

Keywords: adrenal gland; comet assay; DNA damage; loud noise; rat

Aknowledgments

We gratefully acknowledge Mr. Claudio Ghezzani for his excellent technical assistance in the

image analysis. All the authors assure the absence of any conflict of interest regarding the

present article. This work was supported by the Italian Ministry of Research and by the

University of Pisa.

List of abbreviations: Analysis of Variance (ANOVA); Low Melting Point Agarose (LMPA);

Multifactor Analysis of Variance (MANOVA); Normal Melting Point Agarose (NMPA);

Reactive oxygen species (ROS).

2

OUTLINE OF SECTIONS

Abstract

Introduction

Materials and methods

Animals

Experimental procedures

Light microscopy

Evaluation of DNA damage

Results

Discussion

References

Abstract

Loud noise is generally considered as an environmental stressor causing negative effects on acoustic, cardiovascular, nervous and endocrine systems. In this study we investigated the effects of noise exposure on DNA integrity in rat adrenal gland evaluated by the comet assay. The exposure to loud noise (100 dBA) for 12 h caused a significant increase of DNA damage in the adrenal gland. Genetic alterations did not decrease 24 h after the cessation of the stimulus. An imbalance of redox cell status was hypothesized as responsible for the induction and persistence of noise-induced cellular damage.